



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

birds having been secured not uncommonly in the East, and vice versa, does not allow, however, the establishment of a western subspecies.—REGINALD HEBER HOWE, JR., *Longwood, Mass.*

**Notes from Chateaugay Lake, New York.**—During a collecting trip to Chateaugay Lake, Northern Adirondacks, last autumn (Aug. 24 to Sept. 7, 1897) I secured with Mr. G. C. Shattuck, a pair of American Three-toed Woodpeckers (*Picoides americanus*), a species not very uncommon about the lake. One specimen of the Wood Thrush (*Turdus mustelinus*) was also taken. Mr. Shattuck had in previous years taken specimens. This bird was generally found in company with Hermits and Swainson's Thrushes behind the camp where the waste food was thrown. Its occurrence there seems to show that it is found along the western as well as eastern shores of Lake Champlain. Early during my stay I was pretty sure I caught a glimpse of a Philadelphia Vireo (*Vireo philadelphicus*), and after I left Mr. Shattuck secured a specimen.—REGINALD HEBER HOWE, JR., *Longwood, Mass.*

**Ectopistes migratorius, Mimus polyglottos, and Sturnella magna neglecta in Bristol Co., Mass.**—In company with a friend and my brother on August 23, 1889, I was shooting on the mud flats around the reservoir at Norton, Mass. In making a detour of a small inlet, I flushed a Passenger Pigeon from among the low blueberry and bayberry bushes among which I was tramping. The bird alighted in a small white birch near at hand, seeming very unsuspicious, and I shot it. On dissection it proved to be a ♀ young-of-year and was in very good plumage. This is the last record I have of this species. The bird is now mounted and in my collection.

April 30, 1896, a Mockingbird appeared and established himself among the shrubbery in a neighbor's grounds. This was no escaped cage-bird, as his perfect, unfrayed feathers evinced. He was in constant song during his stay, frequently singing half the night when the moon was bright. After enjoying a week of Mockingbird music I was disappointed to find the singer gone, owing to a late driving snow storm, and he did not again appear.

On April 9, of this year (1898), a Western Meadowlark made a visit of a few days in the fields not far from my home. The bird was first seen by an acquaintance, who asked "what bird is it that resembles a Meadowlark in form and color, larger and darker possibly than the Meadowlark but with a wonderfully beautiful song." This bird remained in the same locality for four days, showing no signs of fear and offering an excellent opportunity for one to observe its habits. I was unable to shoot the bird, but identity is beyond all doubt, the song alone being sufficient to remove all queries on that score.—BRADFORD ALEXANDER SCUDDER, *Taunton, Mass.*